

AD-A067 836

VIRGINIA UNIV CHARLOTTESVILLE DEPT OF CHEMISTRY
TETRACARBON METALLOCARBORANES. 6. STEREOCHEMICAL RELATIONSHIPS --ETC (U)
APR 79 J R PIPAL, R N GRIMES

N00014-75-C-0305

F/G 7/3

UNCLASSIFIED

TR-27

NL

OF
AD
A067833



12 LEVEL II

OFFICE OF NAVAL RESEARCH

CONTRACT NO. 15 N00014-75-C-0305

Task No. NR 053-569

9 TECHNICAL REPORT NO. 14 MR-27

6 Tetracarbon Metallocarboranes. 6.
Stereochemical Relationships
Between $\text{Co}_2^{\text{I}}\text{C}_4^{\text{I}}\text{B}_6^{\text{I}}$ Cage Isomers
Formed by Oxidative Fusion. Crystal
Structure of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6^{\text{I}}$
Isomer V.

10 J. Robert Pipal and Russell N. Grimes
Department of Chemistry, University of Virginia
Charlottesville, Virginia 22901

Prepared for Publication

in


Inorganic Chemistry

11 Apr 1979

12 29p.

Reproduction in whole or in part is permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited



401 147

B

79 04 12 029

AD A067836

DDC FILE COPY

Contribution from the Department of Chemistry
University of Virginia
Charlottesville, Virginia 22901

Tetracarbon Metallocarboranes. 6.¹ Stereochemical
Relationships Between $\text{Co}_2\text{C}_4\text{B}_6$ Cage Isomers
Formed by Oxidative Fusion. Crystal Structure of
 $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6$, Isomer V

J. Robert Pipal and Russell N. Grimes*

Abstract. The crystal and molecular structure of the title compound was determined from X-ray diffraction data and found to consist of two pentagonal pyramidal $(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{CH}_3)_2\text{C}_2\text{B}_3\text{H}_3$ units which are partially fused together along their C_2B_3 faces, such that the pairs of carbon atoms on the two pyramidal units are separated by nonbonding distances. The molecule resembles a severely distorted icosahedron with a large opening on one side, and its central cage system is very similar to that previously reported for the monocobalt complex $(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{CH}_3)_4\text{C}_4\text{B}_7\text{H}_6\text{-OC}_2\text{H}_5$. From ^{11}B and ^1H NMR data, the title compound is isostructural with isomer V of the $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2\text{C}_4\text{B}_6\text{H}_{10}$ system, which was previously obtained together with two other isomers (VI and VII) in the oxidative fusion of 1,2,3- $(\eta^5\text{-C}_5\text{H}_5)\text{CoC}_2\text{B}_3\text{H}_7$ in ethanolic KOH. These three isomers adopt different types of cage geometry, as determined from the present study, from an X-ray investigation of VII, and from NMR data on isomer VI. A structure is proposed for isomer VI, and a mechanism is suggested to account for the formation of the three $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2\text{C}_4\text{B}_6\text{H}_{10}$ species from $(\eta^5\text{-C}_5\text{H}_5)\text{CoC}_2\text{B}_3\text{H}_7$ as well as the formation

of a single isomer of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6$ (the title compound) from $(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{CH}_3)_2\text{C}_2\text{B}_3\text{H}_5$. Crystal data for $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6$: space group $I\bar{4}2d$, $Z = 8$; $a = 13.838(2)$, $c = 20.635(5)\text{\AA}$; $V = 3951(1)\text{\AA}^3$; $R = 0.034$ for the 1021 reflections for which $F_o^2 > 3\sigma(F_o^2)$.

| | |
|---------------------------------|---------------------------------------------------|
| ACCESSION for | |
| NTIS | White Section <input checked="" type="checkbox"/> |
| DDC | Buff Section <input type="checkbox"/> |
| UNANNOUNCED | <input type="checkbox"/> |
| JUSTIFICATION _____ | |
| BY _____ | |
| DISTRIBUTION/AVAILABILITY CODES | |
| Dist. | AVAIL. and/or SPECIAL |
| A | |

Introduction

Structural studies of carboranes and metallocarboranes having four carbon atoms in the same polyhedron are proving to be an effective probe of the relationship between electron population and cage geometry, and have revealed a number of unprecedented molecular shapes.¹ Since the formal replacement of a BH by a CH (or C-alkyl) group in a closed polyhedral (closo) system increases the number of valence electrons in the cage skeleton, one can predict from electron-counting arguments² that tetracarbon carboranes will exhibit cage structures that are more open than their dicarbon counterparts; simply put, this is a consequence of the fact that the additional electrons in most cases must occupy antibonding orbitals, thereby producing some sort of cage-opening or distortion. However, this general principle does not tell us what kind of distortion to expect in a particular case, or what the geometry of a particular tetracarbon species will be. This is emphatically the case in 12-vertex, 28-electron cage systems⁴, which contain two more skeletal electrons than do 26-electron icosahedral species such as $C_2B_{10}H_{12}$ and $B_{12}H_{12}^{2-}$. All of the structurally characterized 28-electron cages are distorted from icosahedral geometry, but the nature of the distortion varies widely; as we pointed out recently⁴, at least four different types of cage geometry have been identified among the 12-vertex, 28-electron species that have been structurally characterized.

Most striking are the cases in which isomers produced in the same reaction adopt markedly different polyhedral shapes.^{1,5,6} A recent example¹ is given by the oxidative fusion of the small metallocarborane 1,2,3-(η^5 -C₅H₅)CoC₂B₃H₇ in ethanolic KOH, which produces (η^5 -C₅H₅)CoC₄B₇H₁₁ and three isomers of (η^5 -C₅H₅)₂Co₂C₄B₆H₁₀ (a 12-vertex, 28-electron system). We have previously described the synthetic chemistry together with a crystallographic study of one of the three dicobalt isomers.¹ In this paper we report the structural characterization of a second isomer via its C-tetramethyl derivative, and propose a structure for the third isomer as well as a possible pathway for the formation of all three species.

Experimental Section

Crystals of (C₅H₅)₂Co₂(CH₃)₄C₄B₆H₆ were grown by the vapor diffusion of pentane into a methylene chloride solution of the compound. One of these multifaceted crystals, a rough parallelepiped with dimensions of 0.5 x 0.5 x 0.25 mm, was mounted on a glass fiber in an arbitrary orientation and examined by preliminary precession photographs which indicated high crystal quality. Crystal data: Co₂C₁₈B₆H₂₈; mol wt 427.16; space group I $\bar{4}$ 2d (No. 122); Z = 8; a = 13.838(2), c = 20.635(5) Å; V = 3951(1) Å³; μ (MoK α) = 17.0 cm⁻¹; ρ_c = 1.405 g/cm³; F(000) = 1760. For this crystal the Enraf-Nonius program SEARCH was used to obtain 25

accurately centered reflections which were then used in the program INDEX to obtain an orientation matrix for data collection and to provide approximate cell dimensions. Refined cell dimensions and their estimated standard deviations were determined using these same 25 reflections and the Enraf-Nonius program UNICELL. The mosaicity of the crystal was examined by the ω scan technique and found acceptable. Systematic absences of $h + k + l = 2n + 1$ for $h \ k \ l$, ($k + l = 2n + 1$) on $0 \ k \ l$ and $2h + l \neq 4n$ on $h \ k \ l$, indicate that the space group is either $I4_1md$ or $I\bar{4}2d$. The latter was shown to be correct. For $Z = 8$, this is consistent with the molecular formula assuming 19.0 A^3 per nonhydrogen atom and twofold symmetry for the molecule.

Collection and Reduction of the Data. Diffraction data were collected at 295°K on an Enraf-Nonius four circle CAD-4 diffractometer controlled by a PDP8/M computer, using $\text{MoK}\alpha$ radiation from a highly oriented graphite crystal monochrometer. The θ - 2θ scan technique was used to record the intensities for all reflections for which $1^\circ \leq 2\theta \leq 52^\circ$. Scan widths were calculated from the formula $\text{SW} = A + B \tan \theta$ where A is estimated from the mosaicity of the crystal and B compensates for the increase in the width of the peak due to $\text{K}\alpha_1$ and $\text{K}\alpha_2$ splitting. The values of A and B were 0.60 and 0.35° respectively. This calculated scan angle was extended at each side by 25% for background determination (BG1 and BG2). The net count (NC) was then calculated as $\text{NC} = \text{TOT} - 2(\text{BG1} + \text{BG2})$ where TOT is the estimated peak intensity.

Reflection data were considered insignificant for intensities registering less than ten counts above background on a rapid prescan, and these reflections were rejected automatically by the computer. The intensities of three standard reflections were monitored at intervals of 100 reflections and showed no systematic trends. Raw intensity data were corrected for Lorentz-polarization effects which resulted in a total of 1264 intensities of which 1021 had $F_o^2 \geq 3\sigma(F_o^2)$, where $\sigma(F_o^2)$ was estimated from counting statistics using an ignorance factor⁷ of 0.03. These latter reflections were used in the final refinement of the structural parameters.

Solution and Refinement of the Structure. Initial efforts at solution and refinement were performed assuming $I4_1md$ as the space group. With this choice, solution of the Patterson map for the possible coordinates of the unique cobalt was achieved but many strong peaks in the map were not accounted for. This set of coordinates failed to refine. Various reasonable estimates of the cobalt position consistent with minor symmetry for the molecule (with $Z = 8$) and spectroscopic information (NMR) likewise failed to refine in $I4_1md$. Therefore this initial choice was abandoned in favor of $I\bar{4}2d$. With some difficulty, the Patterson was solved unambiguously for the position of the unique cobalt. The second cobalt in the molecule is related to the first by the crystallographic fourfold roto-inversion operation in $I\bar{4}2d$ carried out twice. Least squares refinement

of the cobalt atom coordinates and thermal parameters reduced the conventional residual R to 0.255. An electron density difference map phased on this refined cobalt yielded ten of the twelve unique remaining nonhydrogen atoms; subsequent maps were used to locate the last two. Isotropic followed by anisotropic refinement lowered R to 0.045 and R_w to 0.069, where R and R_w are defined as $\sum ||F_o| - |F_c|| / \sum |F_o|$ and $(\sum w(|F_o| - |F_c|)^2 / \sum w|F_o|^2)^{1/2}$ respectively.

Several more electron density difference maps were then used to locate the positions of the terminal hydrogens bonded to the boron atoms as well as possible locations for the methyl hydrogens. The former successfully refined with isotropic thermal parameters. The latter did not, and were replaced by hydrogen atoms held fixed at calculated positions 0.95 Å from their respective carbons. Further refinement reduced the residuals to their final values of $R = 0.034$ and $R_w = 0.043$. The estimated standard deviation of an observation of unit weight is 2.301 and the ratio of data to parameters is 7.9. During the last cycle of refinement the largest parameter shift was 0.01 times its estimated error. A structure factor calculation including those data for which $F_o^2 < 3\sigma(F_o^2)$ gave $R = 0.050$. No absorption correction was attempted because of the very irregular multifaceted nature of the crystal. Given the crystal's boxlike shape and its relatively

small absorption coefficient, the error from this source was judged insignificant. A final electron density difference map was featureless.

Full matrix least squares refinement was based on F , and the function minimized was $\sum w(|F_o| - |F_c|)^2$. The weights w were taken as $[2F_o/\sigma(F_o^2)]^2$ where $|F_o|$ and $|F_c|$ are the observed and calculated structure factor amplitudes. The atomic scattering factors for nonhydrogen atoms were taken from Cromer and Waber⁸ and those for hydrogen from Stewart.⁹ The effects of anomalous dispersion were included in F_c using Cromer and Ibers'¹⁰ values of $\Delta f'$ and $\Delta f''$. The computing system and programs are described elsewhere.¹¹ A table of observed and calculated structure factors is available (see paragraph at end of paper regarding supplementary material).



Results and Discussion

Final positional and thermal parameters are given in Table I while Tables II and III contain intramolecular distances and angles. The digits in parentheses in the tables are the estimated standard deviations in the least significant figure quoted and were derived from the inverse matrix in the course of least squares refinement calculations. Tables IV and V list selected mean planes and intermolecular contacts, respectively. Figure 1 is a stereoscopic view of the molecule.

Description of the Structure. The molecule consists of two pyramidal CoC_2B_3 units whose pentagonal faces are partially fused along their respective $\text{B}(4)\text{-B}(5)\text{-B}(6)\text{-C}(2)$ and $\text{B}(4')\text{-B}(5')\text{-B}(6')\text{-C}(2')$ edges (primed and unprimed atoms are related by a crystallographic twofold axis through the molecule). The distances between the pair of framework carbons atoms $\text{C}(2)$, $\text{C}(3)$ and their counterparts $\text{C}(2')$, $\text{C}(3')$ are clearly nonbonding (2.7 Å or greater), so that the cage has a large opening on the side facing the viewer in Figure 1. This geometry is very similar to that of $1,2,3,7,8\text{-(}\eta^5\text{-C}_5\text{H}_5\text{)Co(CH}_3\text{)}_4\text{C}_4\text{B}_7\text{H}_6\text{-OC}_2\text{H}_5$ ⁴ (Figure 2a), an analogue of the present structure in which one $\text{Co(C}_5\text{H}_5\text{)}$ unit has been replaced by $\text{B-OC}_2\text{H}_5$; for comparison, the dihedral angle between the C_2B_3 ring planes is 26.3° in the present structure and 28.5° in the monocobalt species.⁴ Similarly, the $\text{C}(3)\text{-C}(3')$ distance across the open face in the dicobalt species is 2.791(5) Å, while the corresponding vector in the monocobalt structure [$\text{C}(3)\text{-C}(7)$] is 2.854(6) Å. These data indicate that the $\text{Co}_2\text{C}_4\text{B}_6$ framework is slightly less open than the CoC_4B_7 cage. On the other hand, both the $\text{Co}_2\text{C}_4\text{B}_6$ and CoC_4B_7 systems are significantly different from $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ ¹² (Figure 2b), in which the central C-C interaction is bonding [1.53(1) Å]. Since these three species form an

isoelectronic series with 28 skeletal electrons, one can say that formal replacement of one apex BH unit in $(CH_3)_4C_4B_8H_8$ with a $Co(\eta^5-C_5H_5)$ group produces a major structural change, but replacement of the second apex BH has little effect. These findings are important in light of the severe differences that have been observed between $Co_2C_4B_6$ isomers, to be discussed below.

Relationship to the Structures of $(\eta^5-C_5H_5)_2Co_2C_4B_6H_{10}$ Isomers. As described elsewhere¹, the nido complex 1,2,3- $(\eta^5-C_5H_5)CoC_2B_3H_7$ can be deprotonated by treatment with KOH/ethanol or sodium hydride in tetrahydrofuran. The resulting anion, when exposed to air in 10% ethanolic KOH solution, undergoes oxidative fusion² to generate a series of tetracarbon metallo-carboranes as well as other products,¹ in yields of a few percent each.



When the C,C'-dimethyl species, 1,2,3- $(\eta^5-C_5H_5)Co(CH_3)_2C_2B_3H_5$ is treated in identical fashion, the only tetracarbon metallo-carborane obtained is a single isomer of $(\eta^5-C_5H_5)_2Co_2(CH_3)_4C_4B_6H_6$, the compound whose structure we report in this paper. From the 32.1-MHz ¹¹B and 100-MHz ¹H pulse Fourier transform

NMR spectra¹, it is clear that this complex is a C-tetramethyl derivative of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2\text{C}_4\text{B}_6\text{H}_{10}$, isomer V; hence, the gross cage geometry of the tetramethyl species (Figure 1) is assumed to be that of the parent isomer V.

The structures of the two crystallographically characterized isomers, V and VII¹, are clearly different (Figure 3), and in fact constitute different types of nido cages. The geometry of V can be described as a severely distorted icosahedron, while VII resembles a 13-vertex closo polyhedron from which the unique high-coordinate vertex has been removed (this type of cage is also found in $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CH}_3)_4\text{C}_4\text{B}_7\text{H}_8$,⁵ an isoelectronic analogue of VII). From NMR data, to be discussed below, it is highly probable that the remaining isomer, VI, falls into yet another structural class.

The formation of three structurally dissimilar cobalto-carborane isomers at room temperature in the same reaction can only be interpreted in terms of kinetic factors which are dependent on reaction conditions. Whatever the thermodynamically preferred geometry of the $\text{Co}_2\text{C}_4\text{B}_6$ system may be (and it could well be different from any of the three observed isomers V, VI, or VII), the structures of the isolated products no doubt reflect specific pathways by which they are generated from the CoC_2B_3 precursor. In Figure 3 we suggest a scheme for the formation of all three isomers. The initial step probably involves formation of a "quadruple-decker" complex (I) in which two $(\eta^5\text{-C}_5\text{H}_5)\text{CoC}_2\text{B}_3\text{H}_3^{2-}$ ligands are

sandwiched around a central CoH^{4+} group; this process would be precisely analogous to the known synthesis¹³ of $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{CoH}$ from the $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ ion and CoCl_2 in THF. In the present case, the source of the central cobalt ion is doubtless the degradation of the original monocobalt complex in basic media, which is extensive and produces a variety of products.¹ Complex I has not been isolated due to its rapid conversion to other species, but NMR evidence for the existence of its C-tetramethyl derivative has been obtained.¹⁴

Subsequent air-oxidation of I results in oxidative fusion of the two $(\eta^5\text{-C}_5\text{H}_5)\text{CoC}_2\text{B}_3\text{H}_5^{2-}$ ligands to produce the neutral $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2\text{C}_4\text{B}_6\text{H}_{10}$ complexes.¹ In our proposed scheme this occurs through the partially linked intermediate II, which with minor adjustment becomes the observed isomer V; again, this is a process directly analogous to the known conversion,^{13,15} of $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{CoH}$ or $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{FeH}_2$ to $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$. Indeed, as was pointed out earlier, isomer V is a structural counterpart of $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ in which the central C-C interaction is stretched to nonbonding distance.

Intermediate II is also proposed to undergo an alternative type of rearrangement to produce the symmetric intermediate III from which isomers VI and VII are formed as shown. The suggested structure of VI is based on its ^{11}B and ^1H NMR spectra¹, which indicate C_{2v} symmetry with two boron environments in a 2:4 ratio

and equivalent $(C_5H_5)Co$ groups. The area-2 ^{11}B signal appears at very low field (δ 69.1 ppm relative to BF_3 etherate), strongly suggesting the presence of two four-coordinate BH units $[B(5), B(5')]$ located adjacent to both cobalt nuclei.^{16,17} These data are highly restrictive, and while alternative structures cannot be ruled out, a pseudo-icosahedral type cage seems strongly indicated for isomer VI. Mild distortion of the idealized C_{2v} geometry in Figure 3 (perhaps lowering the symmetry to C_2) would not be surprising, but a highly opened framework (such as VII, for example) would be difficult to reconcile with the NMR observations.

The scheme shown in Figure 3 also provides a rationale for the fact that only isomer V of $(\eta^5-C_5H_5)_2Co_2(CH_3)_4C_4B_6H_6$ is obtained when the starting material is $(\eta^5-C_5H_5)Co(CH_3)_2C_2B_3H_5$, as opposed to the three isomers (V, VI, VII) that are formed from the parent (non-methylated) complex. In isomers VI and VII, the framework carbon atoms are in close proximity, requiring the two HC-CH pairs in intermediate III to move toward each other. In isomer V, on the other hand, no such close approach is involved and the two pairs of carbon atoms are in fact well separated. One might then expect that in the C-tetramethylated system, the cage geometry of V would be strongly favored and those of VI and VII hindered, as a consequence of methyl-methyl repulsions.

Conclusions

The structure reported in this paper, taken together with previously established structures,^{1,2,4,12,13} extends our understanding of tetracarbon cobaltacarborane stereochemistry in a significant way: it allows us to consider in some detail the mechanisms of formation and interconversion of the $\text{Co}_2\text{C}_4\text{B}_6$ cage isomers. In general, we appear to have reached a point at which the available structural information in this area can support at least some mechanistic ideas. In other publications we shall attempt to deal with the observed¹⁵ reversible rearrangement of $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ isomers, the formation and stereochemistry of $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8^{2-}$ and $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_9^-$ ions,⁵ the insertion of transition metals into these species to give tetracarbon metallocarboranes of various types,² and related problems.

Acknowledgments. This work was supported in part by the Office of Naval Research and by the National Science Foundation, Grant CHE 76-04491.

Supplementary Material Available: Listing of observed and calculated structure factors (5 pages). Ordering information is given on any current masthead page.

References and Notes

- (1) Part 5: K-S. Wong, J. R. Bowser, J. R. Pipal, and R. N. Grimes, J. Am. Chem. Soc., 100, 5045 (1978).
- (2) For a review, see R.N. Grimes, Acc. Chem. Res., 11, 420 (1978).
- (3) (a) K. Wade, Adv. Inorg. Chem. Radiochem., 18, 1 (1976);
(b) R.W. Rudolph, Acc. Chem. Res., 9, 446 (1976); (c) D.M. P. Mingos, Nature (London), Phys. Sci., 236, 99 (1972).
- (4) J.R. Pipal and R.N. Grimes, J. Am. Chem. Soc., 100, 3083 (1978).
- (5) W.M. Maxwell, R.F. Bryan, and R.N. Grimes, J. Am. Chem. Soc., 99, 4008 (1977).
- (6) W.M. Maxwell, R. Weiss, E. Sinn, and R.N. Grimes, J. Am. Chem. Soc., 99, 4016 (1977).
- (7) P.W.R. Corfield, R.J. Doedens, and J.A. Ibers, Inorg. Chem., 6, 197 (1967).
- (8) D.T. Cromer and J.T. Waber, "International Tables for X-ray Crystallography," Vol. IV, the Kynoch Press, Birmingham, England, 1974.
- (9) R.F. Stewart, E.R. Davidson, and W.T. Simpson, J. Chem. Phys., 42, 3175 (1965).
- (10) D.T. Cromer and J.A. Ibers, ref. 8.
- (11) D.P. Freyberg, G.M. Mockler, and E. Sinn, J. Chem. Soc., Dalton Trans., 447 (1976).
- (12) D.P. Freyberg, R. Weiss, E. Sinn, and R.N. Grimes, Inorg. Chem., 16, 1847 (1977).

- (13) W.M. Maxwell, V.R. Miller, and R.N. Grimes, J. Am. Chem. Soc., 98, 4818 (1976).
- (14) D.F. Finster and R.N. Grimes, unpublished results.
- (15) W.M. Maxwell, V.R. Miller, and R.N. Grimes, Inorg. Chem., 15, 1343 (1976).
- (16) (a) V.R. Miller and R.N. Grimes, J. Am. Chem. Soc., 95, 2830 (1973); (b) W.J. Evans, G.B. Dunks, and M.F. Hawthorne, ibid., 95, 4565 (1973); (c) V.R. Miller and R.N. Grimes, ibid., 97, 4213 (1975).
- (17) Support for this assignment is the fact that (1) no other known tetracarbon metallocarborane has an ^{11}B resonance lower than $\delta \sim 35$ ppm, and (2) no previously characterized tetracarbon metallocarborane has a low-coordinate boron adjacent to more than one metal. Thus, the strikingly different ^{11}B spectrum of VI implies a structural type not previously seen in this class of compounds.
- (18) J.R. Pipal, W.M. Maxwell, and R.N. Grimes. Inorg. Chem., 17, 1447 (1978).

| atom | X | Y | Z | U11 | U22 | U33 | U12 | U13 | U23 |
|-------|------------|------------|------------|-----------|-----------|-----------|-----------|------------|-----------|
| Co | 0.33176(4) | 0.05600(4) | 0.03268(3) | 0.0341(2) | 0.0315(2) | 0.0407(2) | 0.0008(2) | -0.0058(2) | 0.0047(2) |
| C(2) | 0.3941(3) | -0.0445(3) | 0.0899(2) | 0.033(2) | 0.027(2) | 0.036(2) | 0.001(2) | -0.001(2) | 0.005(2) |
| C(3) | 0.4135(3) | 0.0519(3) | 0.1105(2) | 0.034(2) | 0.028(2) | 0.033(2) | 0.004(2) | -0.003(1) | -0.002(2) |
| CH(2) | 0.3393(3) | -0.1166(3) | 0.1304(3) | 0.045(2) | 0.035(2) | 0.069(3) | -0.005(2) | -0.001(3) | 0.012(2) |
| CH(3) | 0.3851(3) | 0.0684(3) | 0.1775(2) | 0.047(2) | 0.048(2) | 0.047(2) | 0.005(2) | 0.004(2) | -0.010(2) |
| CP(1) | 0.2417(4) | 0.1725(4) | 0.0310(4) | 0.052(2) | 0.043(2) | 0.201(6) | 0.021(2) | -0.065(3) | -0.022(4) |
| CP(2) | 0.2542(4) | 0.1319(4) | -0.0358(3) | 0.048(2) | 0.085(3) | 0.102(4) | 0.003(3) | -0.019(3) | 0.051(3) |
| CP(3) | 0.2190(4) | 0.0387(5) | -0.0337(3) | 0.049(2) | 0.085(4) | 0.064(3) | 0.001(3) | -0.024(2) | -0.001(3) |
| CP(4) | 0.1861(4) | 0.0183(4) | 0.0287(3) | 0.042(2) | 0.061(3) | 0.092(4) | -0.001(2) | -0.015(3) | 0.016(3) |
| CP(5) | 0.1979(4) | 0.0974(5) | 0.0650(3) | 0.049(2) | 0.088(4) | 0.067(3) | 0.023(3) | -0.007(3) | -0.003(3) |
| B(4) | 0.4746(3) | 0.1117(3) | 0.0616(3) | 0.035(2) | 0.029(2) | 0.050(2) | 0.001(2) | -0.000(2) | 0.002(2) |
| B(5) | 0.4613(4) | 0.0512(4) | -0.0178(2) | 0.041(2) | 0.048(3) | 0.038(2) | -0.001(2) | 0.002(2) | 0.007(2) |
| B(6) | 0.4181(4) | -0.0665(4) | 0.0132(2) | 0.038(2) | 0.038(2) | 0.040(2) | -0.005(2) | -0.006(2) | -0.010(2) |
| H(4) | 0.490(3) | 0.192(3) | 0.063(2) | 2.1(7) | | | | | |
| H(5) | 0.452(4) | 0.084(4) | -0.069(2) | 5.9(14) | | | | | |
| H(6) | 0.377(3) | -0.122(3) | -0.012(2) | 2.4(8) | | | | | |
| H(21) | 0.334 | -0.176 | 0.107 | 6.0 | | | | | |
| H(22) | 0.372 | -0.127 | 0.170 | 6.0 | | | | | |
| H(23) | 0.276 | -0.092 | 0.139 | 6.0 | | | | | |
| H(31) | 0.404 | 0.154 | 0.182 | 6.0 | | | | | |
| H(32) | 0.317 | 0.084 | 0.182 | 6.0 | | | | | |
| H(33) | 0.416 | 0.051 | 0.210 | 6.0 | | | | | |

THE FORM OF THE ANISOTROPIC THERMAL PARAMETER IS $\exp[-2\pi^2(U_{11}h^2a^2 + U_{22}k^2b^2 + U_{33}l^2c^2 + 2U_{12}hka^*b^* + 2U_{13}hla^*c^* + 2U_{23}klb^*c^*)]$. ^bFor hydrogen atoms, standard isotropic B values are given.

Table II. Interatomic Distances (Å)^a

Bonded Distances

| | | | |
|------------|----------|-------------|----------|
| Co-C(2) | 2.017(3) | C(3)-CM(3) | 1.524(4) |
| Co-C(3) | 1.965(3) | B(4)-B(5) | 1.849(5) |
| Co-B(4) | 2.205(4) | B(4)-B(6') | 1.895(5) |
| Co-B(5) | 2.074(3) | B(4)-H(4) | 1.14(3) |
| Co-B(6) | 2.112(4) | B(5)-B(5') | 1.776(8) |
| Co-CP(1) | 2.039(3) | B(5)-B(6) | 1.848(5) |
| Co-CP(2) | 2.062(4) | B(5)-B(6') | 1.800(5) |
| Co-CP(3) | 2.090(4) | B(5)-H(5) | 1.17(4) |
| Co-CP(4) | 2.084(4) | B(6)-H(6) | 1.09(3) |
| Co-CP(5) | 2.079(4) | CP(1)-CP(2) | 1.500(7) |
| C(2)-C(3) | 1.425(4) | CP(2)-CP(3) | 1.379(6) |
| C(2)-B(4') | 2.122(4) | CP(3)-CP(4) | 1.394(6) |
| C(2)-B(6) | 1.645(4) | CP(4)-CP(5) | 1.385(6) |
| C(2)-CM(2) | 1.507(4) | CP(5)-CP(1) | 1.436(7) |
| C(3)-B(4) | 1.555(4) | | |

Nonbonded Distances

| | | | |
|------------|----------|------------|----------|
| Co-Co' | 4.907(1) | C(2)-C(3') | 2.698(4) |
| C(2)-C(2') | 3.178(6) | C(3)-C(3') | 2.791(5) |

^aAtoms marked with a prime are related to their unmarked counterparts by an inversion axis which bisects the B(5)-B(5') and C(3)-C(3') vectors.

Table III. Selected Bond Angles (deg)

| | | | |
|------------------|----------|-------------------|----------|
| C(2)-Co-C(3) | 41.9(1) | C(2')-B(4)-C(3) | 93.1(2) |
| C(2)-Co-B(6) | 46.9(1) | C(2')-B(4)-B(6') | 47.9(2) |
| C(3)-Co-B(4) | 43.3(1) | B(5)-B(4)-B(6') | 57.4(2) |
| B(4)-Co-B(5) | 51.1(1) | C(3)-B(4)-B(5) | 106.2(2) |
| B(5)-Co-B(6) | 52.4(2) | Co-B(5)-B(4) | 68.1(2) |
| Co-C(2)-C(3) | 67.1(1) | Co-B(5)-B(6) | 64.9(2) |
| Co-C(2)-B(6) | 69.6(2) | B(4)-B(5)-B(6') | 62.6(2) |
| C(3)-C(2)-B(4') | 109.3(2) | B(5')-B(5)-B(6') | 62.3(2) |
| C(3)-C(2)-B(6) | 115.0(3) | B(5')-B(5)-B(6) | 59.5(2) |
| B(4')-C(2)-B(6) | 58.8(2) | B(4)-B(5)-B(6) | 97.2(2) |
| Co-C(2)-CM(2) | 124.5(2) | Co-B(6)-C(2) | 63.5(2) |
| C(3)-C(2)-CM(2) | 123.3(3) | Co-B(6)-B(5) | 62.8(2) |
| B(4')-C(2)-CM(2) | 107.0(2) | C(2)-B(6)-B(4') | 73.3(2) |
| B(6)-C(2)-CM(2) | 120.8(3) | B(4')-B(6)-B(5') | 60.0(2) |
| Co-C(3)-C(2) | 71.0(2) | B(5)-B(6)-B(5') | 58.2(2) |
| Co-C(3)-B(4) | 76.6(2) | C(2)-B(6)-B(5) | 103.6(2) |
| C(2)-C(3)-B(4) | 114.0(3) | CP(2)-CP(1)-CP(5) | 106.2(3) |
| Co-C(3)-CM(3) | 125.7(2) | CP(1)-CP(2)-CP(3) | 106.3(4) |
| C(2)-C(3)-CM(3) | 122.2(3) | CP(2)-CP(3)-CP(4) | 109.5(4) |
| B(4)-C(3)-CM(3) | 123.5(3) | CP(3)-CP(4)-CP(5) | 110.8(4) |
| Co-B(4)-C(3) | 60.1(2) | CP(1)-CP(5)-CP(4) | 107.1(4) |
| Co-B(4)-B(5) | 60.8(2) | | |

Table IV. Selected Intramolecular Planes



| Atom | Deviation | Atom | Deviation |
|-----------------------------------------------------------------------------------------|------------|--------|------------|
| Plane 1: C(2), C(3), B(4), B(5), B(6) 0.9081x - 0.3516y + 0.2274z = 5.5732 | | | |
| C(2) | -0.017 | B(6) | -0.066 |
| C(3) | 0.111 | Co | 1.524 |
| B(4) | -0.137 | CM(2) | 0.131 |
| B(5) | 0.109 | CM(3) | 0.332 |
| Plane 2: C(2'), C(3'), B(4'), B(5'), B(6') 0.9081x - 0.3516y - 0.2274z = 6.9927 | | | |
| Plane 3: CP(1), CP(2), CP(3), CP(4), CP(5) 0.9083x - 0.3337y + 0.2521z = 2.3999 | | | |
| CP(1) | -0.002 | CP(4) | -0.003 |
| CP(2) | 0.000 | CP(5) | 0.003 |
| CP(3) | 0.002 | Co | -1.682 |
| Plane 4: CP(1'), CP(2'), CP(3'), CP(4'), CP(5') 0.9083x - 0.3337y - 0.2521z = 10.169 | | | |
| Planes | Angle, deg | Planes | Angle, deg |
| 1,2 | 26.3 | 2,3 | 27.8 |
| 1,3 | 1.8 | 2,4 | 1.8 |
| 1,4 | 27.8 | 3,4 | 29.2 |




Table V. Intermolecular Non-Hydrogen Contacts ($<3.8 \text{ \AA}$)

| Atom 1 | Atom 2 | Distance, \AA | Relationship |
|--------|--------|------------------------|--------------------|
| CM(2) | CM(2) | 3.699(6) | $x, -1/2-y, 1/4-z$ |
| CM(2) | CP(2) | 3.797(5) | $y, -1/2+x, 1/4+z$ |
| CP(2) | CP(2) | 3.68(1) | $1/2-x, y, -1/4-z$ |
| CP(3) | CP(5) | 3.753(6) | $-y, x, -z$ |


Figure Captions

- Figure 1. Molecular structure of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6$.
Primed atoms are related to the corresponding unprimed atoms by a crystallographic twofold axis bisecting the B(5)-B(5') bond.
- Figure 2. Comparison of the structures of $(\eta^5\text{-C}_5\text{H}_5)\text{Co}(\text{CH}_3)_4\text{C}_4\text{B}_7\text{H}_6\text{-OC}_2\text{H}_5$ (a)⁴ and $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ (b).¹²
- Figure 3. Proposed mechanism of formation of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2\text{C}_4\text{B}_6\text{H}_{10}$ isomers. The structures of isomers V and VII are established; that of VI is proposed from NMR evidence. Species I, II, and III are suggested reaction intermediates. The conversion of II to III can occur via insertion of B(5') between B(5) and B(4) and insertion of B(5) between B(5') and B(4'), with subsequent linkage of B(5') to Co(1) and of B(5) to Co(1'); the other rearrangements depicted are obvious.





OB, BH ● C, CH ◐ CH₃



FORMATION OF $(C_5H_5)_2Co_2C_4B_6H_{10}$ ISOMERS

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|---------------------------------------------------------------------------|
| 1. REPORT NUMBER Technical Report No. 27 | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) Tetracarbon Metallocarboranes. 6. Stereochemical Relationships Between $\text{Co}_2\text{C}_4\text{B}_6$ Cage Isomers Formed by Oxidative Fusion. Crystal Structure of $(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}_2(\text{CH}_3)_4\text{C}_4\text{B}_6\text{H}_6$, Isomer V | | 5. TYPE OF REPORT & PERIOD COVERED Interim |
| 7. AUTHOR(s) J. Robert Pipal and Russell N. Grimes | | 6. PERFORMING ORG. REPORT NUMBER |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Virginia, Department of Chemistry, Charlottesville, Va. 22901 | | 8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0305 |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch, Office of Naval Research Arlington, Virginia 22217 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-569 |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 12. REPORT DATE April, 1979 |
| | | 13. NUMBER OF PAGES 25 |
| | | 15. SECURITY CLASS. (of this report) Unclassified |
| | | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Metallocarboranes Cobaltacarboranes Oxidative cage fusion Tetracarbon metallocarboranes Dimetallocarboranes | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See page 1 | | |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

No.
Copies

Office of Naval Research
800 North Quincy Street
Arlington, Virginia 22217
Attn: Code 472

2

ONR Branch Office
536 S. Clark Street
Chicago, Illinois 60605
Attn: Dr. George Sandoz

1

ONR Branch Office
715 Broadway
New York, New York 10003
Attn: Scientific Dept.

1

ONR Branch Office
1030 East Green Street
Pasadena, California 91106
Attn: Dr. R. J. Marcus

1

ONR Area Office
One Hallidie Plaza, Suite 601
San Francisco, California 94102
Attn: Dr. P. A. Miller

1

ONR Branch Office
Building 114, Section D
666 Summer Street
Boston, Massachusetts 02210
Attn: Dr. L. H. Peebles

1

Director, Naval Research Laboratory
Washington, D.C. 20390
Attn: Code 6100

1

The Assistant Secretary of the Navy
(R, E&S)

Department of the Navy
Room 4E736, Pentagon
Washington, D.C. 20350

1

Commander, Naval Air Systems Command
Department of the Navy
Washington, D.C. 20360
Attn: Code 310C (H. Rosenwasser)

1

Defense Documentation Center
Building 5, Cameron Station
Alexandria, Virginia 22314

12

U.S. Army Research Office
P.O. Box 1211
Research Triangle Park, N.C. 27709
Attn: CRD-AA-IP

1

Naval Ocean Systems Center
San Diego, California 92152
Attn: Mr. Joe McCartney

1

Naval Weapons Center
China Lake, California 93555
Attn: Dr. A. B. Amster
Chemistry Division

1

Naval Civil Engineering Laboratory
Port Hueneme, California 93401
Attn: Dr. R. W. Drisko

1

Professor K. E. Woehler
Department of Physics & Chemistry
Naval Postgraduate School
Monterey, California 93940

1

Dr. A. L. Slafkosky
Scientific Advisor
Commandant of the Marine Corps
(Code RD-1)
Washington, D.C. 20380

1

Office of Naval Research
800 N. Quincy Street
Arlington, Virginia 22217
Attn: Dr. Richard S. Miller

1

Naval Ship Research and Development
Center
Annapolis, Maryland 21401
Attn: Dr. G. Bosmajian
Applied Chemistry Division

1

Naval Ocean Systems Center
San Diego, California 91232
Attn: Dr. S. Yamamoto, Marine
Sciences Division

1

TECHNICAL REPORT DISTRIBUTION LIST, 053

| | <u>No.</u> <u>Copies</u> | | <u>No.</u> <u>Copies</u> |
|--------------------------------------------------------------------------------------------------------------------------|-----------------------------|------------------------------------------------------------------------------------------------------------------------------|-----------------------------|
| Dr. R. N. Grimen University of Virginia Department of Chemistry Charlottesville, Virginia 22901 | 1 | Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401 | 1 |
| Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843 | 1 | Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154 | 1 |
| Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024 | 1 | Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201 | 1 |
| Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401 | 1 | Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802 | 1 |
| Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375 | 1 | Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019 | 1 |
| Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 39716 | 1 | Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214 | 1 |
| Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514 | 1 | Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802 | 1 |
| Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139 | 1 | Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167 | 1 |